



# Acute PresentatiOn of CoLorectal Cancer: an internatiOnal snapshot

**An international, multi-centre study of emergency presentations of colorectal cancer**

30 October 2022  
Version 1.0

## **EuroSurg**

Website: [eurosurg.org](http://eurosurg.org)  
General email: [apollo.eurosurg@gmail.com](mailto:apollo.eurosurg@gmail.com)  
REDCap queries: [apollo.eurosurg@gmail.com](mailto:apollo.eurosurg@gmail.com)  
Twitter: @EuroSurg

## **Key Study Dates**

Study registration opens: 2nd November 2022  
Data collection periods: 16th January 2023  
Last follow-up period ends: 11th September 2023  
ESCP Annual meeting: 20th-23rd September 2023

# Table of Contents

<b>Table of Contents</b>	<b>2</b>
<b>Study Management Group</b>	<b>3</b>
<b>Scientific Advisory Group</b>	<b>3</b>
<b>Collaborative Partners</b>	<b>4</b>
<b>Study Delivery Timeline</b>	<b>5</b>
<b>Executive summary</b>	<b>6</b>
<b>About EuroSurg</b>	<b>7</b>
<b>Introduction</b>	<b>8</b>
<b>Methods</b>	<b>9</b>
1. Study aims	9
2. Study design	9
3. Setting	10
4. Project timeline	10
5. Patient eligibility	11
6. Patient identification	12
7. Outcomes and variables	12
8. Follow up	12
9. Data collection and storage	13
10. Statistical analysis plan	13
11. Local governance and ethical approval	14
12. Quality assurance	15
13. Authorship and mini-teams	15
14. Patient and public involvement	19
15. Expected outputs	19
<b>Acknowledgements</b>	<b>20</b>
<b>Appendix A: APOLLO Study Case Report Form</b>	<b>21</b>
<b>Appendix B: Data Dictionary</b>	<b>23</b>
<b>Appendix C: National Lead Survey</b>	<b>30</b>
<b>Appendix D: Clavien-Dindo Grading of Surgical Complications</b>	<b>31</b>
<b>Appendix E: American Society of Anaesthesiologists (ASA) Classification</b>	<b>33</b>
<b>Appendix F: How to make your centre a success</b>	<b>34</b>
<b>References</b>	<b>36</b>

## Study Management Group

Name	Training stage	Twitter
Niall Brindl	Resident Surgeon, Heidelberg	@BrindlNiall
Daoud Chaudhry	Foundation Doctor (FY1), Stoke-On-Trent	@da_chaudhry
Kassandra Gressmann	4th year medical student, Dublin	
Amie Hilder	3rd year medical student, Melbourne	@amie_hilder
Esther Matey	Foundation Doctor (FY1), Scotland	
Irène Santos	6th year medical student, Porto	@IreneSantos_P
Adam Turňa	4th year medical student, London	@AdamTurna2
Laura Kehoe	5th year medical student, Dublin	
Mustafa Ege Seker	5th year medical student, Istanbul	@MEgeSeker
Noor Essa	5th year medical student, Dublin	
Mafalda Sampaio Alves	5th year medical student, Porto	@sampalvmaf
Muhammed Elhadi	Medical doctor, Tripoli	@ElhadiMuhammed
Chris Varghese	Trainee intern, University of Auckland	@chrsvarghese98
William Xu	Trainee intern, University of Auckland	@williamxu_98

## Scientific Advisory Group

Name	Position	Twitter
Ian Bissett	Professor and consultant colorectal surgeon	@ian_bissett
Sue Blackwell	PPI representative at ESCP	@littlemissileo
Stephen Chapman	Research fellow and surgical trainee	@SJ_Chapman
Ruth Blanco Colino	Surgical trainee	@ruthbc93
Eloy Espin	Professor and consultant colorectal surgeon	@eloiespin
James Glasbey	NIHR Doctoral Research Fellow, Birmingham, UK	@DrJamesGlasbey
Susan Moug	Professor and consultant colorectal surgeon	@susanmoug
Dion Morton	Professor and consultant colorectal surgeon	
Dmitri Nepogodiev	Public Health Registrar	@dnepo
Francesco Pata	Consultant colorectal surgeon	@drfrancescopata
Gianluca Pellino	Consultant colorectal surgeon	@GianlucaPellino
Peter Pockney	Professor and consultant colorectal surgeon	@PPockney
Alessandro Sgrò	Surgical trainee	@AlessandroSgr1
Wal Baraza	Consultant colorectal surgeon	

## Collaborative Partners



**STARSurg Collaborative**

<https://starsurg.org/>  
[starsurguk@gmail.com](mailto:starsurguk@gmail.com)  
Twitter: @STARSurgUK



**itSurg Collaborative**

[www.itsurg.org](http://www.itsurg.org)  
[itsurg.group@gmail.com](mailto:itsurg.group@gmail.com)  
Twitter: @it\_surg



**PTSurg Collaborative**

[www.ptsurg.org](http://www.ptsurg.org)  
[info@ptsurg.org](mailto:info@ptsurg.org)  
Twitter: @pt\_surg



**SIGMA Collaborative**

[www.SIGMA-studies.org](http://www.SIGMA-studies.org)  
[team@SIGMA-studies.org](mailto:team@SIGMA-studies.org)  
Twitter: @sigmastudies



**TASMAN Collaborative**

<https://anzsurgsocs.org/tasman/>  
[tasmananz@gmail.com](mailto:tasmananz@gmail.com)  
Twitter: @TASMANCollab

## Study Delivery Timeline

Dates	Description
<b>Study set up</b>	
Sep 2022	Provisional proposal released at EuroSurg session, ESCP 2022, Dublin
1 Nov 2022	Final protocol release
Oct - Dec 2022	National committees established Patient involvement to prioritise outcomes
Nov - Jan 2022	Application for ethical approval at local centres Application for REDCap Logins
16 Jan 2023	First Collaborator REDCap Accounts Generated (then on a rolling twice weekly basis for all new collaborators – Tuesdays and Fridays)
<b>Data collection</b>	
16 January 2023 - 13 June 2023	Any period of 6 weeks within this 5 month period for patient recruitment. Additional periods can be completed for a total of 12/18 weeks
11 September 2023	Last day of 90-day follow-up
<b>Data validation and analysis</b>	
30 September 2023	Data validation
15 October 2023	REDCap Database Locked, Final Data Submission Deadline
30 October 2023	Deadline for data validation
November 2023	Data analysis and write up
Sep 2023	Preliminary results presented at ESCP 2023 (first half of cohort)

## Executive summary

**Primary aim:** Describe the variation in the operative and non-operative management of emergency presentations of colon and rectal cancer in an international cohort.

**Secondary aims:**

- Identify risk factors for mortality (intraoperatively, at 30-days and at 90-days) and ostomy rates (at 30- and 90-days) in patients deemed for active management (i.e., not for palliative management) to develop a risk prediction model
- Validate risk criteria of large bowel obstruction in patients with previously known colorectal cancer undergoing neoadjuvant chemotherapy or awaiting elective surgery

**Who?**

- Patients aged 18 years and above presenting to the hospital acutely with colorectal cancer (CRC) for malignant large bowel obstruction (LBO), perforation, CRC-related haemorrhage or other reasons within the data collection periods. Both those managed with and without surgery will be included
- Patients with localised and metastatic disease will be included
- Patients with known colorectal cancer diagnoses will be included if they present acutely (e.g., with disease progression)
  
- Patients presenting acutely for the side effects of chemotherapy/radiotherapy of known cancers will be excluded

**What?**

Data will be collected on patients' presenting status and symptoms, patient management strategies, and intraoperative and postoperative outcomes.

**When?**

Prospectively over 2023 in consecutive 6-week data collection blocks between January and June with 90-day follow-up till September.

## About EuroSurg

The EuroSurg collaborative is an international research group led by students and surgical trainees. Founded at the European Society of Coloproctology (ESCP) 2015 meeting, it has since expanded rapidly with active members in Czech Republic, France, Germany, Italy, Netherlands, Spain, Turkey, Portugal, Ireland, the United Kingdom, Sweden, Norway, Greece, Hungary, Latvia, South Africa, New Zealand and Australia.<sup>1</sup> In our most recent study COMPASS<sup>2,3</sup> which explored the use of peritoneal drains after elective and emergency colorectal surgery, over 20 different countries were involved. Previous EuroSurg projects can be found at <http://eurosurg.org/> and include EuroSurg-1,<sup>4</sup> IMAGINE,<sup>5-7</sup> COMPASS,<sup>3</sup> and CASCADE.<sup>8</sup>

The model for trainee-led research collaboratives was pioneered in the UK by local networks of trainee surgeons. These networks have been successful in delivering major surgical research initiatives, including multicentre cohort studies and randomised controlled trials (RCTs). The feasibility of students conducting similar projects was first demonstrated by the Student Audit & Research in Surgery (STARSurG) collaborative, which has delivered several national cohort studies in the UK.<sup>9,10</sup>

Collaboration across international surgical communities produces transferable results which may inform the design of future RCTs and changes in clinical practice. Through participating in a EuroSurg project, students will acquire essential skills in surgical audit and research methodology. EuroSurg's authorship policy designates all student and trainee collaborators as PubMed-citable "Collaborators". An example of this authorship model can be seen here.<sup>3,5</sup>

## Introduction

Colorectal cancer (CRC) presents as an emergency in as many as a third of patients,<sup>11,12</sup> 80% of the time with obstruction and, less commonly, perforation and haemorrhage.<sup>13</sup> Emergency surgery for CRC is associated with mortality in 15-34%, morbidity in 32-64%,<sup>14</sup> higher ostomy rates, and poorer health related quality of life,<sup>14</sup> Patients presenting emergently with CRC also tend towards more deranged physiology and advanced tumour biology.<sup>13,15</sup>

Existing guidelines on the optimal management of such cases are based on predominantly low grade evidence,<sup>16</sup> and current series remain small. Global practices in the management of emergency CRC remain unquantified. There are various options for patients presenting acutely with colorectal cancer, such as primary resection with anastomosis, colonic stenting as a bridge to elective surgery, primary resection with an end colostomy, or defunctioning stomas. Furthermore, with the rising use of neoadjuvant chemotherapy for colonic cancer,<sup>17</sup> management pathways have become increasingly complex. Accurate operative and anastomosis risk prediction in acutely presenting colorectal cancer may help guide clinical decision making.

A significant number of patients presenting with malignant large bowel obstruction also have disseminated disease.<sup>18</sup> For these patients, self-expandable metal stents may offer a lower risk of permanent stoma compared to traditional surgical options, but also may result in perforations.<sup>19,20</sup> The use of stents worldwide remains unquantified.

There is impetus for an international snapshot audit of emergency presentations of CRC to capture prospective outcome data and describe the variation in management worldwide. These data would be useful to externally validate ongoing trials, and identify new research gaps to power new trials.

**APOLLO** (Acute PresentatiOn of CoLorectal Cancer: an internatiOnal snapshot) is an international, multi-centre, prospective observational study which will address this need and aims to describe the operative and non-operative management of emergency presentations of colon and rectal cancer in an international cohort.

# Methods


## 1. Study aims

The primary aim of the **APOLLO** (Acute PresentatiOn of CoLorectal Cancer: an internatiOnal snapshot) study is to describe the operative and non-operative management of emergency presentations of colon and rectal cancer in an international cohort. Secondary aims will be to describe 30-day and 90-day management outcomes, identify the risk factors for intraoperative, 30-day, and 90-day mortality and ostomy rates in patients deemed for active management (i.e., not for palliative management), and develop a mortality and ostomy risk prediction model for patients undergoing active management for colorectal cancer. This study will also aim to validate risk criteria of large bowel obstruction in patients with previously known colorectal cancer undergoing neoadjuvant chemotherapy or awaiting elective surgery

## 2. Study design

**APOLLO** is an international, multi-centre, prospective observational study of acutely (i.e. unplanned and non-elective presentation to hospital for urgent or emergency reasons) presenting colorectal cancer. This study will adapt the student- and trainee-led collaborative research model. 'Mini-teams' of collaborators will participate at each hospital, with a range of members including medical students, junior doctors, trainees, registrars, and supervising consultants (**Figure 1**). It is compulsory to have a consultant supervisor who is able to guide and advise how you may register the study at your hospital, and what approvals will be required. **It will also be compulsory to have 1 general surgical trainee as a member of your mini-team.**

**Figure 1: Mini-team structure**



At each hospital, one mini-team per period will collect data on eligible patients admitted over a consecutive 6-week period. Patients should be included if their hospital admission started (defined as 'date of hospital admission') within the time period during the data collection periods as specified above.

### 3. Setting

**APOLLO** is open to any secondary or tertiary hospitals across the world with a general or colorectal surgery department performing major colorectal cancer surgery. All participating centres will be required to register the study according to local regulations, evidence of which must be uploaded onto REDCap prior to the commencement of data collection from each respective site. As this is an audit of practice, no change to normal patient management is required.

In the UK, **APOLLO** has been designated an audit of practice. Internationally, individual study investigators are responsible for ensuring the correct approvals have been achieved prior to commencing data collection. At each hospital, one mini-team per period will collect data on eligible patients admitted over a consecutive 6-week period. Following the conclusion of the study, it is recommended that mini-teams at each centre present the study findings to their hospital's surgery departments in order to close the audit loop.

### 4. Project timeline

Collaborators at each participating centre will prospectively collect data for all eligible patients over minimum 6-week periods from January to June 2023. Sites can contribute in multiple 6-week periods, providing they are **not** overlapping.

There will be a rolling start date from 16 January, 2023 to 1 May, 2023. The last date a patient can be recruited is the 13 June. All 90-day follow-up should be completed by 11 September, 2023. Recruitment dates may be extended at the discretion of the steering committee, allowing for extensions if there are delays in site initiation.

## 5. Patient eligibility

**Summary:** Consecutive adult patients ( $\geq 18$  years of age) presenting acutely (i.e. unplanned and non-elective presentation to hospital for urgent or emergency reasons) for symptoms of known or unknown colorectal cancer assessed by hospital surgical teams. Patients should be included regardless of operative or non-operative management, and curative or palliative intent.

Inclusion criteria	
Age	18 years and above
Patient pathology	<p>Patients admitted to the hospital acutely with <b>primary colon AND/OR rectal adenocarcinoma</b> and referred to general/colorectal surgical departments</p> <p>This includes:</p> <ul style="list-style-type: none"> <li>• Patients who are operated on for curative/palliative treatment of colorectal cancer</li> <li>• Patients who are referred to general surgery for assessment (irrespective of if they proceed to surgery)</li> <li>• New diagnoses of colorectal cancer</li> </ul>
Extent of cancer	All patients including extra-abdominal metastatic, intra-abdominal metastatic and non-metastatic disease
Known cancer	<p>Patients presenting for the first time with colorectal cancer <b>AND/OR</b></p> <p>Known colorectal cancer diagnosis with progression of disease. i.e. obstruction of known colorectal cancer</p>
Patient presentation	Symptomatic presentation of colorectal cancer i.e. large bowel obstruction, haemorrhage, perforation
Exclusion criteria	
Presenting for side effects of cancer treatment	Patients presenting acutely for the side effects of chemotherapy/radiotherapy of known cancers are excluded
Patients with non-colorectal primary cancers	Non-primary colorectal cancers that have metastasised to the colon and rectum (e.g., melanoma or lymphoma). Primary neuroendocrine tumours, gastrointestinal lymphomas, gastrointestinal stromal tumours, and primary colorectal squamous cell carcinomas should also be excluded.
Previously included in dataset	Patients should only be included once in APOLLO

***All eligible patients must be included to avoid selection bias.***

## 6. Patient identification

Teams should collect data on consecutive eligible patients at their hospital during the data collection period. Strategies to identify consecutive patients could include:

- Daily review of theatre lists
- Daily review of handover meetings/sheets and ward lists
- Daily review of theatre logbooks
- Daily review of new inpatient referrals to surgical team
- Daily on call review of patients undergoing emergency surgery
- Colorectal cancer multidisciplinary team meetings

## 7. Outcomes and variables

**Primary outcome:** 90-day mortality since day of presentation

**Secondary outcomes include:**

- Inpatient mortality, 30-day mortality since day of presentation
- Operative 30- and 90-day mortality rates
- Rates of primary anastomosis
- 30- and 90-day stoma formation and reversal rates
- Rate and grade of surgical complications
- Rates of stenting in those with left-sided LBO, rates of stenting in those treated with palliative intent <sup>21,22</sup>, rate of colonic stenting complications
- Rate of representation for patients managed nonoperatively
- Rates of radiological assessment on admission
- Proportion of patients presenting acutely with large bowel obstruction with known colorectal cancer who had high risk criteria for obstruction (as defined by the FOXTroT obstruction criteria) at index assessment

**Audit standard outcomes:**

- All patients with suspected LBO should have a contrast-enhanced CT <sup>23</sup>
- <20% 90-day mortality after emergency colorectal cancer surgery (grade b) (consistent with NBOCA recommendations)

**Variables**

- Covariates are detailed in **Appendix B**

## 8. Follow up

Patients will be followed up for 90-days after presentation. No change in routine follow up should take place. Collaborators should be proactive in identifying follow-up data, but this should be done within the limits of routine follow up.

Strategies for follow-up include:

- Regularly reviewing patient notes to identify in-hospital complications
- Participating in daily ward rounds and doctor reviews
- Reviewing clinic notes and clinic letters, if seen in clinic by 30 days
- Checking electronic systems and handover lists for re-admissions
- Checking for emergency department re-attendances

## 9. Data collection and storage

Data will be collected and stored online via the Research Electronic Data Capture (REDCap) web application,<sup>24,25</sup> hosted and managed by the Birmingham Surgical Trials Consortium (BiSTC) REDCap system hosted at the University of Birmingham, United Kingdom.

REDCap allows collaborators to enter and store data in a secure system. It is widely used by academic institutions throughout Europe and all storage of web-based information by this system is encrypted and compliant with HIPAA-Security Guidelines in the United States.

The security of the study database system is governed by the policies of the University of Birmingham. Data management and data security within the BiSTC REDCap will abide by the requirements of the General Data Protection Regulations (GDPR) and any subsequent amendments. Collaborators will be given secure REDCap project server login details, allowing secure data storage on the REDCap system.

**No** patient data will be uploaded or stored on the REDCap database without prior local permissions.

All data should be handled in accordance with local data governance policies, and all paper copies of any data should be destroyed as confidential waste within the centre once uploaded to REDCap.

It will not be possible to store patient identification numbers (hospital numbers) on REDCap. A unique 'REDCap ID' will be generated by the system for each patient. If needed, you may keep a local cross-reference of hospital numbers and REDCap IDs. This should be kept in a secure, encrypted spreadsheet on a hospital, password-protected computer. No patient identifiable data will be uploaded or stored on the REDCap database.

**One** REDCap login will be issued per collaborator and **only** that person may use the login. If you experience problems, please email [apollo.eurosurg@gmail.com](mailto:apollo.eurosurg@gmail.com).

Data collected during the **APOLLO** study can be used for future analyses at the Study Management Group's discretion.

## 10. Statistical analysis plan

The primary descriptive analysis will be investigating emergency CRC presentation and their respective management.

- Presenting symptoms — rates of obstruction, perforation and bleeding
- Management approaches — resection, primary anastomosis, end-stoma formation, stents, palliation
- Outcomes — rates of mortality, ostomy rates

Planned secondary analyses include:

- A risk prediction tool in emergency CRC surgery with curative intent for 30- and 90-day mortality will be created per the TRIPOD statement to guide management in emergency colorectal cancer surgery if the recruitment numbers allow (LBO ± perforation/others). This will allow colorectal surgeons to preoperatively predict short term morbidity and mortality and determine which patients may instead benefit from a lower-risk bridging stent before elective surgery.
- A risk prediction tool for whether to anastomosis or not (preoperative) to aid decision making

For model development, a mixed effects logistic regression model will be built with candidate variables selected based on clinical plausibility. Relative model fits will be assessed using Akaike information criterion (AIC) to obtain the most parsimonious model. Area under the receiver operating characteristic curve (AUROC) will be used to assess discrimination. The linearity assumption for logistic regression will be assessed by plotting. Validation of the model will be performed as the methods of Bonsdorff et al <sup>26</sup>. Statistical analysis will be completed using R (R Foundation Statistical Program).

## 11. Local governance and ethical approval

It is compulsory to have a consultant supervisor who is able to guide and advise how you may register the study at your hospital, and what approvals will be required. These must be added to the REDCap database as evidence of approvals. You may also seek advice from your local audit department, or get in touch with the [apollo.eurosurg@gmail.com](mailto:apollo.eurosurg@gmail.com) for further advice. **It will also be compulsory to have a minimum of 1 general surgical registrar as members of your mini-team when possible (2 recommended if possible)**. If there is only one or no general surgical registrar available, team structures will be approved on a case-by-case basis.

In the UK, the study may be registered as a clinical audit or “service evaluation” to assess mortality after emergency colorectal cancer surgery:

*<20% 90-day mortality after emergency colorectal cancer surgery (grade b)  
(consistent with NBOCA recommendations)*

When registering the study, the following points should be made clear:

- All data collected will measure current practice

- No changes to normal patient pathways/ treatment will be made
- This is an international audit

UK collaborators should also seek their NHS trust's Caldicott Guardian's approval to submit data to the REDCap system.

*Collaborators should complete the mandatory data governance e-learning module which will be made available on the online project hub: <http://www.eurosurg.org>*

## 12. Quality assurance

**Design:** This protocol was written with guidance from an international expert cross-speciality advisory group and with the contribution of patient representatives. A data dictionary will be developed to help collaborators in collecting data and patient inclusion. E-learning materials will be available on our website (eurosurg.org).

**Data completeness:** Following data collection, only data sets with >95% data completeness will be accepted for pooled analysis. To emphasise the importance of data completeness to collaborators, centres with >5% missing data points will be excluded from the study and collaborators from those centres withdrawn from the published list of citable collaborators. See further details under 'Authorship and miniteams'.

**Validation:** Data validation will be performed to ensure all consecutive cases were recruited. Data validators may be either a final year student or a qualified doctor who were not involved in the initial data collection.

The validator will be required to validate a minimum of 1x 6-week period for the centre for which they are acting as the data validator. The validator will require to review every case to confirm case ascertainment and subsequently check a random 20% of collected data points to ensure accurate data collection.

After completing validation, the validator will send a summary of how many records were reviewed and error rates to the study management group. There are two components of validation:

### **I. Case ascertainment:**

The validator will independently identify all patients eligible for inclusion over the one 6-week study period. The target for data ascertainment is >95%.

### **II. Data collection**

The validator will independently collect data for the key data fields relating to risk-adjustment and outcome measures (see **Appendix A**).

Any conflicts with the data originally submitted by the relevant mini-team will be resolved by discussion between the validator and the mini-team. The target for accuracy of collected data is >98%.

### 13. Authorship and mini-teams


Medical students will take the lead in disseminating and delivering this study, which is supported by national committees of medical students, trainees and consultant/attending surgeons.

In accordance with National Research Collaborative (NRC) authorship guidelines <sup>27</sup>, all research outputs from **APOLLO** will be listed under a single corporate authorship (“EuroSurg Collaborative”). All collaborators will be listed as PubMed-citable collaborators within the EuroSurg Collaborative in accordance with the roles defined below (so long as the minimum requirements for authorship are achieved):

**To be credited with authorship, all collaborators must provide a valid ORCID identifier (<https://orcid.org/register>). Authorship lists for all papers will be made from these.**

Mini-teams for the **APOLLO** study can consist of up to 12 members, including a minimum of 1 supervising consultant, and 1 general surgical registrar (*for small centres with no trainees, this requirement will be waived*). Any of the 12 mini-team members can be the **Hospital Lead** and will be responsible for maintaining communication with the EuroSurg Steering Committee.

*Tip: We recommend that 6 of the 9 other members are allocated to data collection over a 6-week consecutive period, and 3 of the 9 other members are allocated to collecting 30- and 90-day follow-up data.*



Collaborator role descriptions are as follows:

1. **Local supervising consultant/attending:** provide guidance for approval processes, facilitate communication within the hospital, and mentor and facilitate medical students, junior doctors and registrars in conducting the study at your local site. They have overall responsibility for the site governance registration and should support data collection. Only one person can fulfil this role. Minimum requirements for authorship include:
  - Sponsorship of local study registration, and responsibility to ensure local collaborators act in accordance with local governance guidelines.
  - Successful completion of data collection at a centre which meets the criteria for inclusion within the **APOLLO** dataset.
  - Facilitation of local result presentation and support of appropriate local interventions.

Sponsorship through the audit approval / project registration process by a consultant does not constitute authorship, nor does inclusion of a consultants' patients alone in the audit serve sufficient for authorship.

2. **Hospital lead:** this role can be fulfilled by a medical student, junior doctor, trainee or the consultant supervisor/PI (as above). Prior experience in collaborative research is recommended but not essential. Additional support can be sought from EuroSurg. They will be the single lead point of contact for data collection at each site and will liaise with the local PI and EuroSurg. You must be responsive to communication from the PI, governance bodies, and EuroSurg.
  - Primary person responsible in obtaining local approvals for conduct of the **APOLLO** Study (e.g. registration of the audit, seeking permission to upload data to REDCap).
  - Successful completion of data collection at a centre which meets the criteria for inclusion within the **APOLLO** dataset.
  - A hospital lead guide can be found in **Appendix D**.
3. **General Surgical Registrar/Resident/Trainee:** a minimum of 1 member of the mini-team is required to be general surgical registrar. It is recommended 2 general surgical registrars/trainees are included in your team if possible. Their role will be:
  - Providing advice around clinical details around colorectal cancer, and acute presentations
  - Liaising with multidisciplinary teams to aid complete case ascertainment
4. **Local collaborators:** a team of up to 9 data collectors per specialty, per centre). Minimum requirements for authorship on **APOLLO** publications include:
  - Compliance with local audit approval processes and data governance policies.
  - Active involvement in data collection over at least one data collection period at a centre which meets the criteria for inclusion within the **APOLLO** dataset.
  - Collaboration with the hospital leads to ensure that the audit results are reported back to the audit office / clinical teams.

- 5. Local validators:** A final year medical student or junior doctor not involved with data collection whose role is to ensure adequate **data ascertainment** and **data collection accuracy** (see 12. Quality assurance)
  - The validator will be a 6-week data collection period at their local centre to validate. Data validation will occur following completion of data collection (including follow-up).
  - After completing validation, the validator will send a summary of how many records were reviewed and error rates to the study management group.
  
- 6. National network committees:** A core group of medical students and surgical trainees in each participating country responsible for study dissemination, protocol translation and supporting students to correctly register and run the study at each participating centre.
  
- 7. Writing Group:** A group of medical students, junior doctors and external advisory board members responsible for the overall scientific content, data analysis, and preparation of individual research manuscripts.
  
- 8. Study management group:** A core group of medical students and junior doctors who have overall responsibility for protocol design, project coordination, and data handling
  
- 9. Data Management and Statistical Analysis group:** A sub team who take overall responsibility for the quality assurance of data analysis and statistical analysis plans.
  
- 10. External Advisory Group:** A panel of cross-disciplinary field experts who are able to ensure contextual and scientific relevance of the protocol design, data fields and data interpretation.

#### **Criteria for site inclusion within APOLLO**

- Successful in obtaining all relevant local approvals for conduct of the **APOLLO** Study
- Have completed the site survey
- Successful data collection of at least one eligible patient per period for each site
- Individual sites must also ensure
  1. They obtain **>95% data completeness** for all required fields
  2. All data has been uploaded by the specified database closure deadline

Should these criteria not be met, the contributing mini-team and any data they contribute may not be included in the final study, and they may be removed from any authorship lists. You are advised to get in touch with us as soon as possible so we may support you with ensuring your site is able to successfully collect data towards the **APOLLO** Study.

For guidance relating to mini-team setup and audit registration, please contact your local supervising consultant or local lead. If you would be interested in signing up as a PI for a new centre not currently involved, or for any general enquiries regarding the protocol, please contact us via email ([apollo.eurosurg@gmail.com](mailto:apollo.eurosurg@gmail.com)) or Twitter (@EuroSurg)

## 14. Patient and public involvement

To better understand the outcomes important to patients living with colorectal cancer we asked the opinion of patients, aiming to clarify what matters to them most in the acute period. Patients were invited to fill out an anonymous survey through social media.

The main questions concerned their experience with presenting to the hospital in an emergency with colorectal cancer, with an emphasis on which outcomes after surgery were most important to them. Furthermore, an emphasis was placed on understanding what information was presented to them regarding treatment options in the acute setting.

Patient and public involvement in this study will be reported according to the short form of the Guidance for Reporting Involvement of Patients and the Public (GRIPP2) reporting checklist (<https://www.bmj.com/content/358/bmj.j3453>).

## 15. Expected outputs

Unit level data for comparison will be fed back to collaborators to support local service improvement (upon request). This project will be submitted for presentation at national and international conferences. Manuscript(s) will be prepared following close of the project.

## Acknowledgements

We thank the **Birmingham Surgical Trials Consortium (BiSTC)** for kindly hosting **APOLLO** on their REDCap servers.

Website: <https://www.birmingham.ac.uk/research/activity/mds/trials/BiSTC/index.aspx>



We thank the **European Society of Coloproctology (ESCP)** for their kind support.

Website: <https://www.escp.eu.com/>



# Appendix A: APOLLO Study Case Report Form

## APOLLO Case Report Form (CRF)

Use with Appendix (Data Dictionary) to help data collection.

Section 1: Baseline data fields		Date of presentation		REDCap unique ID
Age	Sex <input type="checkbox"/> M <input type="checkbox"/> F	ASA Grade <input type="checkbox"/> I <input type="checkbox"/> II <input type="checkbox"/> III <input type="checkbox"/> IV <input type="checkbox"/> V	Weight (kg)	Data collection date range
Clinical frailty score <input type="checkbox"/> 1. Very fit <input type="checkbox"/> 2. Well <input type="checkbox"/> 3. Managing well <input type="checkbox"/> 4. Vulnerable <input type="checkbox"/> 5. Mildly frail <input type="checkbox"/> 6. Moderately frail <input type="checkbox"/> 7. Severely frail <input type="checkbox"/> 8. Very severely frail <input type="checkbox"/> 9. Terminally ill	Charlson comorbidity index	<input type="checkbox"/> MI <input type="checkbox"/> CHF <input type="checkbox"/> PVD <input type="checkbox"/> CVA/ TIA <input type="checkbox"/> Dementia <input type="checkbox"/> COPD <input type="checkbox"/> Peptic Ulcer Disease <input type="checkbox"/> Hemiplegia <input type="checkbox"/> Leukaemia <input type="checkbox"/> Lymphoma <input type="checkbox"/> AIDS		
		Diabetes mellitus	<input type="checkbox"/> Diet controlled <input type="checkbox"/> Uncomplicated <input type="checkbox"/> End-organ damage	
		Liver disease	<input type="checkbox"/> Mild <input type="checkbox"/> Moderate <input type="checkbox"/> Severe	
		CKD	<input type="checkbox"/> eGFR < 60 <input type="checkbox"/> Dialysis <input type="checkbox"/> Post kidney transplant <input type="checkbox"/> uraemia	
		Solid tumour	<input type="checkbox"/> Localised <input type="checkbox"/> Metastatic <input type="checkbox"/> None of the Above	
Smoking Status <input type="checkbox"/> Current <input type="checkbox"/> Stopped smoking <6 weeks <input type="checkbox"/> Stopped smoking 6 weeks - 1 year <input type="checkbox"/> Stopped smoking >1 year <input type="checkbox"/> Never smoked <input type="checkbox"/> Unknown	History of abdominal surgery <input type="checkbox"/> Yes <input type="checkbox"/> No	Including this operation, number of operations in the 30-days prior to procedure	<input type="checkbox"/> 1 operation <input type="checkbox"/> 2 operations <input type="checkbox"/> >2 operations	
Section 2: Disease details				
Known cancer <input type="checkbox"/> Yes - date of diagnosis dd/mm/YYYY <input type="checkbox"/> No	If known cancer, status of known cancer	Reason for acute presentation		<input type="checkbox"/> Bowel obstruction <input type="checkbox"/> Haemorrhage <input type="checkbox"/> Anaemia <input type="checkbox"/> Perforation <input type="checkbox"/> Other (specify)
Location of tumour		<input type="checkbox"/> Caecum <input type="checkbox"/> Ascending colon <input type="checkbox"/> Hepatic Flexure <input type="checkbox"/> Transverse colon <input type="checkbox"/> Splenic flexure <input type="checkbox"/> Descending colon <input type="checkbox"/> Sigmoid or rectosigmoid <input type="checkbox"/> Rectum <input type="checkbox"/> Synchronous → branching		
FIT status in past year <input type="checkbox"/> Positive <input type="checkbox"/> Negative <input type="checkbox"/> Unknown	Colonic investigation in past year If yes:	<input type="checkbox"/> Colonoscopy <input type="checkbox"/> CT colonography <input type="checkbox"/> CT colonoscopy minimal preparation <input type="checkbox"/> Malignancy diagnosed <input type="checkbox"/> Polyp resected <input type="checkbox"/> Polyp not resected <input type="checkbox"/> Nil findings		
Section 3: Radiological and endoscopic investigations				
Radiological assessment Yes: <input type="checkbox"/> Non-contrast CT, <input type="checkbox"/> Contrast CT, <input type="checkbox"/> MRI rectum, <input type="checkbox"/> Abdominal XR, <input type="checkbox"/> Other (specify)	<input type="checkbox"/> No	Radiological features	<input type="checkbox"/> Circumferential tumour <input type="checkbox"/> Stricture tumour <input type="checkbox"/> Obstructing tumour	
Endoscopic assessment <input type="checkbox"/> Yes	<input type="checkbox"/> No	Endoscopic findings	<input type="checkbox"/> Unable to pass scope <input type="checkbox"/> Circumferential tumour <input type="checkbox"/> Stricture tumour <input type="checkbox"/> Ulcerating tumour <input type="checkbox"/> Polypoid tumour. <input type="checkbox"/> Obstructing tumour	
T-stage <input type="checkbox"/> T1 - Invades submucosa <input type="checkbox"/> T2 - Invades muscularis propria <input type="checkbox"/> T3 - Beyond muscularis propria <input type="checkbox"/> T4 - Adjacent organs/peritoneum	N-stage	<input type="checkbox"/> NX (cannot be assessed) <input type="checkbox"/> N0. <input type="checkbox"/> N1 (1-3 nodes) <input type="checkbox"/> N2 (4+ nodes)	M-stage	<input type="checkbox"/> MX (cannot be assessed) <input type="checkbox"/> M0 (no distant metastases) <input type="checkbox"/> M1 (distant metastases)
Mismatch repair gene status		<input type="checkbox"/> Proficient <input type="checkbox"/> Deficient <input type="checkbox"/> Unknown		
If <b>known cancer</b> , were CT/PET assessments done at the <i>previous</i> time of diagnosis	<input type="checkbox"/> Yes <input type="checkbox"/> No	Tick all radiological features that apply for the across:	<input type="checkbox"/> Circumferential tumour on radiology; <input type="checkbox"/> Stricture tumour on radiology	
If <b>known cancer</b> , was endoscopy done at the <i>previous</i> time of diagnosis?	<input type="checkbox"/> Yes <input type="checkbox"/> No	Tick all endoscopic features that apply for the across:	<input type="checkbox"/> Unable to pass endoscope beyond tumour <input type="checkbox"/> Stricture disease on endoscopy	
Section 4: Management				
Intent of treatment	<input type="checkbox"/> Curative (treat the cancer) <input type="checkbox"/> Palliative (symptomatic treatment i.e. relief of obstruction without intention to attempt cure of cancer)			
Immediate management at this presentation (tick all that apply)	<input type="checkbox"/> Surgical <input type="checkbox"/> Procedural <input type="checkbox"/> Non-operative <input type="checkbox"/> Other (specify)			
<input type="checkbox"/> Resection: <input type="checkbox"/> Right <input type="checkbox"/> Left <input type="checkbox"/> Subtotal <input type="checkbox"/> Total colectomy <input type="checkbox"/> Hartmann's procedure <input type="checkbox"/> Stoma: <input type="checkbox"/> Loop <input type="checkbox"/> End <input type="checkbox"/> Primary anastomosis <input type="checkbox"/> Colonic metallic stent <input type="checkbox"/> Transanal drainage tube				
Section 5: Surgical variables if undertaken				
Operative urgency <input type="checkbox"/> Immediate <input type="checkbox"/> Urgent <input type="checkbox"/> Expedited <input type="checkbox"/> Elective	Intra-operative contamination <input type="checkbox"/> Clean-contaminated <input type="checkbox"/> Contaminated <input type="checkbox"/> Dirty	Operative approach <input type="checkbox"/> Open <input type="checkbox"/> Laparoscopic <input type="checkbox"/> Laparoscopic-assisted <input type="checkbox"/> Laparoscopic converted to open <input type="checkbox"/> Robotic <input type="checkbox"/> Robotic converted to open		
Intraoperative complications <input type="checkbox"/> None <input type="checkbox"/> Vascular injury <input type="checkbox"/> Bowel injury <input type="checkbox"/> Injury other organs or structures	Surgeon skill level <input type="checkbox"/> Consultant colorectal surgeon <input type="checkbox"/> Emergency general surgeon/non-colorectal surgeon <input type="checkbox"/> Surgical resident/registrar <input type="checkbox"/> Other _____	Critical care unit admission <input type="checkbox"/> Yes - planned <input type="checkbox"/> Yes - unplanned <input type="checkbox"/> No		
Section 6: Endoscopic procedure variables if stenting undertaken for obstructing tumour				
Endoscopic urgency <input type="checkbox"/> Immediate <input type="checkbox"/> Urgent <input type="checkbox"/> Expedited <input type="checkbox"/> Elective	Endoscopic complications <input type="checkbox"/> None <input type="checkbox"/> Technical failure and inadequate decompression <input type="checkbox"/> Bowel perforation <input type="checkbox"/> Procedure abandoned	Proceduralist skill level <input type="checkbox"/> Consultant colorectal surgeon <input type="checkbox"/> Consultant gastroenterologist <input type="checkbox"/> Surgical/gastroenterology resident/registrar <input type="checkbox"/> Other _____		

Section 7: 30-day follow up										
Location	<input type="checkbox"/> Inpatient admission dd/mm/yy		<input type="checkbox"/> Discharged from index admission dd/mm/yy		Highest 30-day Clavien-Dindo complication grade (if surgery)			<input type="checkbox"/> None <input type="checkbox"/> I <input type="checkbox"/> II <input type="checkbox"/> IIIa/IIIb <input type="checkbox"/> IVa/IVb <input type="checkbox"/> V		
Section 8: 90-day follow up										
Location	<input type="checkbox"/> Inpatient admission dd/mm/yy <input type="checkbox"/> Discharged from index admission dd/mm/yy <input type="checkbox"/> Dead dd/mm/yy		Highest 90-day Clavien-Dindo (if surgery)		<input type="checkbox"/> None <input type="checkbox"/> I <input type="checkbox"/> II <input type="checkbox"/> IIIa/IIIb <input type="checkbox"/> IVa/IVb <input type="checkbox"/> V		Pathological results	<input type="checkbox"/> pR0 <input type="checkbox"/> pR1 <input type="checkbox"/> pR2	Started neoadjuvant chemotherapy	<input type="checkbox"/> Yes dd/mm/yy <input type="checkbox"/> No
Late complications of colonic stents?	<input type="checkbox"/> Yes dd/mm/yy Specify: <input type="checkbox"/> Migration <input type="checkbox"/> Obstruction <input type="checkbox"/> Other	<input type="checkbox"/> No <input type="checkbox"/> NA	Stoma reversed within 90 days		<input type="checkbox"/> Yes dd/mm/yy <input type="checkbox"/> No <input type="checkbox"/> NA	Elective colorectal resection completed if no primary surgery		<input type="checkbox"/> Yes dd/mm/yy <input type="checkbox"/> No <input type="checkbox"/> NA	Anastomotic leak (if surgery)	<input type="checkbox"/> Yes dd/mm/yy <input type="checkbox"/> No
Readmission within 90 days	<input type="checkbox"/> Yes dd/mm/yy <input type="checkbox"/> No	Reason for readmission:		<input type="checkbox"/> Bowel obstruction <input type="checkbox"/> Haemorrhage <input type="checkbox"/> Anaemia <input type="checkbox"/> Perforation <input type="checkbox"/> Other		Discharge date of readmission		dd/mm/yy <input type="checkbox"/> NA	Reoperation within 90 days	<input type="checkbox"/> Yes dd/mm/yy <input type="checkbox"/> No

## Appendix B: Data Dictionary

Baseline data fields	Required data (definition/comment)
Data collection period	Start date - end date
Age	Years (at time of presentation)
Sex	Male/Female
American Society of Anesthesiologists (ASA) physical status	I, II, III, IV, V
Body Mass Index	(BMI) Height, weight, BMI (calculator)
Smoking	Current - cigarette Current - vaping Previous (stopped smoking <6 weeks) - cigarette Previous (stopped smoking <6 weeks) - vaping Previous (stopped smoking 6 weeks – 1 year) - cigarette Previous (stopped smoking 6 weeks – 1 year) - vaping Previous (stopped smoking >1 year) - cigarette Previous (stopped smoking >1 year) - vaping Never smoked
Charlson comorbidity index	Myocardial Infarction (MI) Congestive Heart Failure (CHF) Peripheral Vascular Disease (PVD) Cerebrovascular Accident (CVA) or Transient Ischaemic Attack (TIA) Dementia Chronic Obstructive Pulmonary Disease (COPD) Peptic Ulcer Disease Liver Disease (mild - chronic hepatitis or cirrhosis w/o portal hypertension, moderate - cirrhosis with portal hypertension but no variceal bleeding hx, severe - cirrhosis with portal hypertension with variceal bleeding hx) Diabetes Mellitus (none or diet controlled, uncomplicated, end-organ damage) Chronic Kidney Disease (CKD) Estimated Glomerular Filtration Rate (eGFR) <60/ml/min/1.73m <sup>2</sup> , dialysis or post kidney transplant, or uraemia. Hemiplegia Solid tumour (localised, metastatic) Leukaemia Lymphoma AIDS None of the Above
Clinical Frailty Score <sup>28</sup>	1. <b>Very fit.</b> Robust, active, energetic, well-motivated, and fit; these people commonly exercise regularly and are in the most fit group for their age.

	<ol style="list-style-type: none"> <li>2. <b>Well.</b> People who have no active disease symptoms but are less fit than people in category 1. Often, they exercise or are very active occasionally, e.g., seasonally.</li> <li>3. <b>Managing well.</b> People whose medical problems are well controlled but are not regularly active beyond routine walking.</li> <li>4. <b>Vulnerable.</b> While not dependent on others for daily help, often symptoms limit activities. A common complaint is being “slowed up”, and/or being tired during the day.</li> <li>5. <b>Mildly frail.</b> These people often have more evident slowing and need help in high order instrumental activities of daily living (finances, transportation, heavy housework, medications). Typically, mild frailty progressively impairs shopping and walking outside alone, meal preparation and housework.</li> <li>6. <b>Moderately frail.</b> People need help with all outside activities and with keeping house. Inside, they often have problems with stairs and need help with bathing and might need minimal assistance (cuing, standby) with dressing.</li> <li>7. <b>Severely frail.</b> Completely dependent for personal care, from whatever cause (physical or cognitive). Even so, they seem stable and not at high risk of dying (within ~ 6 months).</li> <li>8. <b>Very severely frail.</b> Completely dependent, approaching the end of life. Typically, they could not recover even from a minor illness.</li> <li>9. <b>Terminally ill.</b> Approaching the end of life. This category applies to people with a life expectancy &lt;6 months, who are not otherwise evidently frail.</li> </ol>
History of abdominal surgery	Yes/no
Including this operation, number of operations this patient has had in the 30 day period prior to this procedure	1 operation 2 operations >2 operations
<b>Disease details</b>	
Date of acute presentation	dd/mm/YYYY
Known cancer	Yes - known cancer (date of diagnosis) No - first presentation of colorectal cancer

Reason for acute presentation (tick all that apply)	Bowel obstruction Haemorrhage/anaemia Perforation Other (specify)
<i>If known cancer</i> Status of known cancer	Awaiting elective surgery Undergoing neoadjuvant chemotherapy/radiotherapy before surgery Non-surgical management (including palliative) Other (specify)
Location of tumour	Caecum Ascending colon Hepatic Flexure Transverse colon Splenic flexure Descending colon Sigmoid or rectosigmoid Rectum Synchronous → fill second checkbox
FOBT (faecal occult blood test) status in past year	Positive Negative Unknown
FIT (faecal immunochemical test) status in past year	Positive Negative Unknown
Colonic investigation in past year	Colonoscopy/CT colonography/CT colonoscopy minimal preparation  If yes: Yes - malignancy diagnosed Yes - polyp resected Yes - polyp not resected Nil findings
<b>Radiological and endoscopic investigations</b>	
<b>Radiological assessment</b> completed on <i>this</i> presentation (tick all that apply)	Yes - non-contrast CT Yes - contrast CT Yes - MRI rectum Yes - abdominal X-ray Yes - other (specify) No
T-stage (preferably on contrast staging CT)	T1 - Invades submucosa T2 - Invades muscularis propria T3 - Beyond muscularis propria T4 - Adjacent organs or peritoneum
N-stage	NX (cannot be assessed)

	N0 N1 (1-3 nodes) N2 (4+ nodes)
M-stage	MX (cannot be assessed) M0 (no distant metastases) M1 (distant metastases)
Radiological features (tick all that apply)	Circumferential tumour Strictureing tumour Obstructing tumour
Mismatch repair gene status	Deficient Proficient Unknown
<b>Endoscopic assessment</b> completed at <i>this</i> assessment	Yes No
Endoscopic findings (tick all that apply)	Unable to pass scope Circumferential tumour Strictureing tumour Ulcerating tumour Polypoid tumour Obstructing tumour
<i>If known cancer</i> , were CT/PET assessments done at the <i>previous</i> time of diagnosis?	Yes/No
Tick all radiological features that apply for the above	Circumferential tumour on radiology; Strictureing tumour on radiology
<i>If known cancer</i> , was endoscopy done at the <i>previous</i> time of diagnosis?	Yes/No
Tick all endoscopic features that apply for the above	Unable to pass endoscope beyond tumour; Strictureing disease on endoscopy
<b>Management</b>	
Is the intent of treatment curative or palliative?	Curative (treat the cancer) Palliative (symptomatic treatment i.e. relief of obstruction without intention to attempt cure of cancer)
Immediate management at <i>this</i> presentation of colorectal cancer	Surgical <ul style="list-style-type: none"> <li>● Resection - Yes/no <ul style="list-style-type: none"> <li>○ Right/left/subtotal/total colectomy/Hartmann's procedure</li> </ul> </li> <li>● Stoma - yes (loop)/ yes (end)/no</li> <li>● Primary anastomosis - yes/no</li> </ul> Procedural <ul style="list-style-type: none"> <li>● Colonic metallic stent</li> <li>● Transanal drainage tube</li> </ul>

	Non-operative management Other (specify)
<b>Surgical variables</b> <i>If surgery undertaken</i>	
<i>If surgical management</i> What is the urgency of surgical intervention - NCEPOD classification	Immediate Urgent Expedited Elective
<i>Intraoperative contamination</i>	<b>Clean-Contaminated:</b> (Gastrointestinal (GI) and genitourinary (GU) tracts entered but no gross contamination). <b>Contaminated:</b> (GI or GU tracts entered with gross spillage or major break in sterile technique). <b>Dirty:</b> (There is already contamination prior to operation, e.g. faeces or bile).
<i>If surgical management</i> What was the operative approach	<b>Open</b> (performed exclusively using instruments inserted into the abdomen through a surgical incision). <b>Laparoscopic</b> (performed exclusively using instruments inserted in to the abdomen through small ports) <b>Laparoscopic-assisted</b> (laparoscopic surgery in which an incision is enlarged to deliver a specimen or to insert a gloved hand into the abdomen). <b>Laparoscopic converted to open</b> (surgery planned to be performed laparoscopically but for unforeseen reasons the decision was made to change to an open approach) <b>Robotic</b> (robot-assisted surgery with no conversion to either laparoscopic or open approaches). <b>Robotic converted to open</b> (surgery planned to be performed robotically but for unforeseen reasons the decision was made to change to an open approach).
Intraoperative complications	None / Vascular injury / Bowel injury (e.g. duodenum) / Injury to other organs or structures (e.g. ureter) / Intraoperative mortality
Surgeon skill level	Consultant colorectal surgeon Emergency general surgeon/non-colorectal surgeon Surgical resident/registrar Other
Critical care unit admission	Yes - planned Yes - unplanned No
<b>Endoscopic procedure variables</b> <i>If stenting undertaken for obstructing tumour</i>	
<i>If endoscopic management</i> What is the urgency of surgical intervention	Immediate Urgent Expedited

NCEPOD classification	Elective
Endoscopic complications	None / Technical failure and inadequate decompression / Bowel perforation / / procedure abandoned
Proceduralist skill level	Consultant colorectal surgeon Consultant gastroenterologist Surgical/gastroenterology resident/registrar Other (specify)
<b>30-day follow up</b>	
Outcome	Inpatient Dead ( <i>specify days from index admission that death occurred</i> ) Discharged (date of discharge from index admission)
Highest 30-day Clavien-Dindo complication grade ( <i>if surgery</i> )	None I II IIIa/IIIb IVa/IVb V
<b>90-day follow up</b>	
Outcome	Inpatient Dead ( <i>specify days from index admission that death occurred</i> ) Discharged (date of discharge from index admission)
Highest 90-day Clavien-Dindo complication grade ( <i>if surgery</i> )	None I II IIIa/IIIb IVa/IVb V
Pathological results	pR0, pR1, pR2
Started neoadjuvant chemotherapy	Yes (specify date) No
<i>If colonic stent was placed, were there late complications within 90 days?</i>	Yes (specify date) Stent migration / Stent obstruction / Other (specify) No
<i>If a stoma was formed, was the stoma reversed within 90 days?</i>	Yes (specify date) No
<i>If primary resection <u>did not</u> occur at index admission, was elective surgery for resection completed within 90 days?</i>	Yes (specify date) No

Did the patient readmission within 90 days for reasons other than surgery?	Yes (specify date) No
Reason for representation	Bowel obstruction Haemorrhage/anaemia Perforation Other (specify)
Date of discharge from readmission	Specify date - to calculate <i>days alive and out of hospital</i>
<i>If the patient had surgery at any point within 90 days, did an anastomotic leak occur?</i>	Yes (specify date) No
LOS of index admission	Days
Reoperation within 90 days	Yes (specify date) No

## Appendix C: National Lead Survey

Data field	Required data
Is there a national colorectal cancer screening programme in place in your country?	Yes/No
If yes, what is the method of screening	Faecal immunochemical test (FIT), sigmoidoscopy, colonoscopy, CT colonography, other stool test, other test
If yes, what is the inclusion criteria for screening?	Age [...] to [...] Other criteria <i>Please provide reference</i>
If FIT is used, what was the cut-off threshold for a positive test?	In screening: µg Hb/g  In symptomatics: µg Hb/g
Centre volume	Number of colorectal resections/year
Hospital beds	Number of beds
Number of colorectal surgical consultants	
Is there a colorectal multidisciplinary team meeting (MDT)?	Yes (if yes - number of colorectal cancer diagnoses passed through MDT over the entire 6-week period) No
Is your centre offering neoadjuvant chemotherapy for colon cancer	Yes

## Appendix D: Clavien-Dindo Grading of Surgical Complications

Adverse postoperative events may be divided up into treatment failures, sequelae and complications.

- **Failure of treatment** occurs when the original surgery fails to achieve its intended benefits; for example, persistent pain following laparoscopic cholecystectomy or tumour recurrence following cancer surgery.
- **Sequelae** are the recognised consequences of a given procedure; for example, gut malabsorption following a large small bowel resection or immune deficiency following splenectomy.
- **Complication:** Any deviation from the normal postoperative course that has an adverse effect on the patient and is not either a treatment failure or sequel.

In the Clavien-Dindo classification,<sup>29</sup> the factor determining the severity of a complication is the treatment required. Consequently, a given complication may be graded differently depending on how it has been managed. For example, an anastomotic leak may be managed just with antibiotics if it is contained (grade II) or it may require re-operation under anaesthetic (grade III).

Some other considerations:

- Intra-operative complications are not considered unless they have an adverse effect on the patient post-operatively. The only exception to this is **intra-operative death**; this is classified as grade V.
- **All postoperative adverse events** are included, even when there is no direct relationship to the surgery.
- **All adverse events within the follow-up** period (30 days) are included, even if they occur following discharge.
- **Diagnostic procedures** are not included. For example, a diagnostic oesophagoduodenoscopy (OGD) to look for a source of bleeding without any intervention would not be considered a complication, but a therapeutic OGD with clipping of a bleeding vessel would be considered a grade III complication. Since negative exploratory laparotomies are considered to be diagnostic procedures, they should not be recorded as complications.

Grade	Definition
Grade I	Any deviation from the normal postoperative course not requiring surgical, endoscopic, or radiological intervention. This includes the need for certain drugs (e.g. antiemetics, antipyretics, analgesics, diuretics, and electrolytes), treatment with physiotherapy, and wound infections that are opened at the bedside

Grade II	Complications requiring drug treatments other than those allowed for Grade I complications; this includes blood transfusion and total parenteral nutrition (TPN)
Grade III	Complications requiring surgical, endoscopic or radiological intervention <ul style="list-style-type: none"> <li>• Grade IIIa - intervention not under general anaesthetic</li> <li>• Grade IIIb - intervention under general anaesthetic</li> </ul>
Grade IV	Life-threatening complications; this includes CNS complications (e.g. brain haemorrhage, ischaemic stroke, subarachnoid haemorrhage) which require intensive care, but excludes transient ischaemic attacks (TIAs) <ul style="list-style-type: none"> <li>• Grade IVa - single-organ dysfunction (including dialysis)</li> <li>• Grade IVb - multi-organ dysfunction</li> </ul>
Grade V	Death of the patient

## Appendix E: American Society of Anaesthesiologists (ASA) Classification

<b>ASA Class</b>	<b>Definition</b>
I	A normally healthy patient
II	A patient with mild systemic disease
III	A patient with severe systemic disease
IV	A patient with severe systemic disease that is a constant threat to life
V	A moribund patient who is not expected to survive without the operation
VI	A declared brain-dead patient whose organs are being removed for donor purposes

## Appendix F: How to make your centre a success

Further details can be found at the **APOLLO study hub** and on the **APOLLO local lead guide**

As a hospital lead, you are responsible for coordinating the APOLLO study at your hospital. This document will guide you through some of the steps you can take to make the project a success at your centre. For further information, please contact your national committee.

### 1. Publicise the project

At your local medical school and hospital site, aim to advertise the APOLLO study as widely as possible. This may include the following approaches:

- Promotional posters
- Public announcements in lectures and hospital/university meetings
- Social media, including creating a Facebook group or Twitter page
- Emails to local surgical departments, societies, and associations
- Emails to individuals who may be interested in the project

Gaining early interest/momentum in the local project delivery will ensure maximum participation from students and trainees/residents. It is also essential to gain the support of a local surgical consultant/attending at each hospital. Strategies to achieve this may include:


- Presenting the APOLLO protocol at local hospital/university meetings
- Arranging a meeting with the head of department to discuss the protocol

### 2. Recruit collaborators to ‘mini-teams’

As a hospital lead, you are responsible for creating and managing “mini-teams” of students and trainees/residents at each hospital. Mini-teams for the **APOLLO** study can consist of up to 12 members, including a minimum of 1 supervising consultant, and a minimum of 1 general surgical registrars (2 recommended).

*Tip: We recommend that 6 of the 9 other members are allocated to data collection over a 6-week consecutive period, and 3 of the 9 other members are allocated to collecting 30- and 90-day follow-up data.*

Collaborators at each participating centre will prospectively collect data for all eligible patients over minimum 6-week periods from January to June 2023. Sites can contribute in multiple 6-week periods, providing they are not overlapping.



The same mini-team may collect data across multiple different data collection periods, if they so wish. However, each mini-team must collect data across different data collection periods, i.e. no patients may be included more than once in the REDCap database.

### 3. Get audit or ethical approval early

All countries involved in the APOLLO study will have different requirements for study or research approval. Speak to your national committee for further advice on the requirements in your country, or to seek advice about local guidelines. They may be able to provide details required for registration and help you to facilitate local approvals. This process can take a long time so it is important to start this as early as possible!

### 4. Prepare for the Study Launch

Once you have created mini-teams at your local hospitals and gained study approval, you are almost ready to start collecting data for the APOLLO study. There are some final important steps to ensure all collaborators are ready to begin data collection. This includes:

- Arrange a local meeting with all collaborators to discuss the study and answer questions
- Ensure that all mini-teams are complete and know where/when they are due to collect data
- Ensure all collaborators have a copy of the full protocol
- Ensure that all collaborators are registered with the REDCap system

## References

1. EuroSurg Collaborative. EuroSurg: a new European student-driven research network in surgery. *Colorectal Dis.* 2016 Feb;18(2):214–5.
2. EuroSurg Collaborative. Management of COMPLicAted intra-abdominal collectionS after colorectal Surgery (COMPASS): protocol for a multicentre, observational, prospective international study of drain placement practices in colorectal surgery. *Colorectal Dis.* 2020 Dec;22(12):2315–21.
3. EuroSurg Collaborative. Intraperitoneal drain placement and outcomes after elective colorectal surgery: international matched, prospective, cohort study. *Br J Surg.* 2022 May 16;109(6):520–9.
4. EuroSurg Collaborative. Body mass index and complications following major gastrointestinal surgery: a prospective, international cohort study and meta-analysis. *Colorectal Dis.* 2018 Aug;20(8):O215–25.
5. EuroSurg Collaborative. Safety and efficacy of non-steroidal anti-inflammatory drugs to reduce ileus after colorectal surgery. *Br J Surg.* 2020 Jan;107(2):e161–9.
6. EuroSurg Collaborative. Safety of hospital discharge before return of bowel function after elective colorectal surgery. *Br J Surg.* 2020 Apr;107(5):552–9.
7. EuroSurg Collaborative. Timing of nasogastric tube insertion and the risk of postoperative pneumonia: an international, prospective cohort study. *Colorectal Dis.* 2020 Dec;22(12):2288–97.
8. Student Audit and Research in Surgery (STARSurg) Collaborative and European Surgical (EuroSurg) Collaborative. CArdiovaSCulAr outcomes after major abDominal surgEry: study protocol for a multicentre, observational, prospective, international audit of postoperative cardiac complications after major abdominal surgery. *Br J Anaesth.* 2022 May;128(5):e324–7.
9. STARSurg Collaborative and TASMAN Collaborative. Evaluation of prognostic risk models for postoperative pulmonary complications in adult patients undergoing major abdominal surgery: a systematic review and international external validation cohort study. *Lancet Digit Health.* 2022 Jul;4(7):e520–31.
10. STARSurg Collaborative. Perioperative intravenous contrast administration and the incidence of acute kidney injury after major gastrointestinal surgery: prospective, multicentre cohort study. *Br J Surg.* 2020 Jul;107(8):1023–32.
11. Barnett A, Cedar A, Siddiqui F, Herzig D, Fowlkes E, Thomas CR Jr. Colorectal cancer emergencies. *J Gastrointest Cancer.* 2013 Jun;44(2):132–42.
12. Gunnarsson H, Holm T, Ekholm A, Olsson LI. Emergency presentation of colon cancer is most frequent during summer. *Colorectal Dis.* 2011 Jun;13(6):663–8.
13. Baer C, Menon R, Bastawrous S, Bastawrous A. Emergency Presentations of Colorectal Cancer. *Surg Clin North Am.* 2017 Jun;97(3):529–45.
14. van Hooft JE, Bemelman WA, Oldenburg B, Marinelli AW, Lutke Holzik MF, Grubben MJ, et al. Colonic stenting versus emergency surgery for acute left-sided malignant colonic obstruction: a multicentre randomised trial. *Lancet Oncol.* 2011

Apr;12(4):344–52.

15. Wong SKC, Jalaludin BB, Morgan MJ, Berthelsen AS, Morgan A, Gatenby AH, et al. Tumor pathology and long-term survival in emergency colorectal cancer. *Dis Colon Rectum*. 2008 Feb;51(2):223–30.
16. Ansaloni L, Andersson RE, Bazzoli F, Catena F, Cennamo V, Di Saverio S, et al. Guidelines in the management of obstructing cancer of the left colon: consensus conference of the world society of emergency surgery (WSES) and peritoneum and surgery (PnS) society [Internet]. Vol. 5, *World Journal of Emergency Surgery*. 2010. Available from: <http://dx.doi.org/10.1186/1749-7922-5-29>
17. Seymour MT, Morton D. FOxTROT: an international randomised controlled trial in 1052 patients (pts) evaluating neoadjuvant chemotherapy (NAC) for colon cancer. *J Clin Orthod*. 2019 May 20;37(15\_suppl):3504–3504.
18. Sebastian S, Johnston S, Geoghegan T, Torreggiani W, Buckley M. Pooled analysis of the efficacy and safety of self-expanding metal stenting in malignant colorectal obstruction. *Am J Gastroenterol*. 2004 Oct;99(10):2051–7.
19. Veld J, Umans D, van Halsema E, Amelung F, Fernandes D, Lee MS, et al. Self-expandable metal stent (SEMS) placement or emergency surgery as palliative treatment for obstructive colorectal cancer: A systematic review and meta-analysis. *Crit Rev Oncol Hematol*. 2020 Nov;155:103110.
20. Hill J, Kay C, Morton D, Magill L, Handley K, Gray RG. CREST: Randomised phase III study of stenting as a bridge to surgery in obstructing colorectal cancer—Results of the UK ColoRectal Endoscopic Stenting Trial (CREST). *J Clin Orthod*. 2016 May 20;34(15\_suppl):3507–3507.
21. N. B, M. K, P. H. Colorectal cancer: Summary of NICE guidance. *BMJ*. 2020;368:m461.
22. Colorectal Cancer. National Institute for Health and Care Excellence (UK);
23. Moran B, Cunningham C, Singh T, Sagar P, Bradbury J, Geh I, et al. Association of Coloproctology of Great Britain & Ireland (ACPGBI): Guidelines for the Management of Cancer of the Colon, Rectum and Anus (2017) - Surgical Management. *Colorectal Dis*. 2017 Jul;19 Suppl 1:18–36.
24. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009 Apr;42(2):377–81.
25. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O’Neal L, et al. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform*. 2019 Jul;95:103208.
26. Bonsdorff A, Ghorbani P, Helanterä I, Tarvainen T, Kontio T, Belfrage H, et al. Development and external validation of DISPAIR fistula risk score for clinically relevant postoperative pancreatic fistula risk after distal pancreatectomy. *Br J Surg* [Internet]. 2022 Aug 19; Available from: <http://dx.doi.org/10.1093/bjs/znac266>
27. National Research Collaborative & Association of Surgeons in Training Collaborative Consensus Group. Recognising contributions to work in research collaboratives: Guidelines for standardising reporting of authorship in collaborative research. *Int J Surg*.

2018 Apr;52:355–60.

28. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005 Aug 30;173(5):489–95.
29. Clavien PA, Barkun J, de Oliveira ML, Vauthey JN, Dindo D, Schulick RD, et al. The Clavien-Dindo classification of surgical complications: five-year experience. *Ann Surg*. 2009 Aug;250(2):187–96.